

References

- 5 Bakalyar, H.A., and Reed, R.R. (1990). Identification of
a specialized adenylyl cyclase that may mediate odorant
detection. *Science* 250, 1403-1406.
- 10 Baldwin, J.M. (1993). The probable arrangement of the
helices in G protein-coupled receptors. *The EMBO Journal*
12(4), 1693-1703.
- 15 Bean, N.J. (1982). Modulation of agonistic behavior by
the dual olfactory system in male mice. *Physiol. Behav.*
29, 433-437.
- 20 Ben Arie, N., Lancet, D., Taylor, C., Khen, M., Walker,
N., Ledbetter, D.H., Carrozzo, R., Patel, K., Sheer, D.,
Lehrach, H., and North, M.A. (1994). Olfactory receptor
gene cluster on human chromosome 17: possible
duplication of an ancestral receptor repertoire. *Hum.*
Mol. Gen. 3, 229-235.
- Bertmar, G. (1981). Evolution of vomeronasal organs in
vertebrates. *Evolution* 35, 359-366.
- 25 Brady, G., Barbara, M., and Iscove, N.N. (1990).
Representative *in vitro* cDNA amplification from
individual hemopoietic cells and colonies. *Methods in*
Mol. Cell. Biol. 2, 17-25.
- 30 Broadwell, R.D. (1975). Olfactory relationships of the
Telencephalon and Diencephalon in the rabbit. I. An
autoradiographic study of the efferent connections of the
main and accessory olfactory bulbs *J. Comp. Neurol.* 163,
329-346.
- 35

Buck, L., and Axel, R. (1991). A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. *Cell* 65, 175-187.

- 5 Halpern, M., Shapiro, L.S., and Jia, C. (1995). Differential localization of G proteins in the opossum vomeronasal system. *Brain Research* 677, 157-161.

- 10 Chess, A., Simon, I., Cedar, H., and Axel, R. (1994). Allelic inactivation regulates olfactory receptor gene expression. *Cell* 78, 823-834.

- 15 Clancy, A.N., Coquelin, A., Macrides, F., Gorski, R.A., and Nobel, E.P. (1984). Sexual behavior and aggression in male mice: involvement of the vomeronasal system. *J. Neurosci.* 4, 2222-2229.

- 20 Dhallan, R.S., Yau, K.W., Schrader, K.A., and Reed, R.R. (1990). Primary structure and functional expression of a cyclic nucleotide-activated channel from olfactory neurons. *Nature* 347, 184-187.

- 25 Eisthen, H.L. (1992). Phylogeny of the vomeronasal system and of receptor cell types in the olfactory and vomeronasal epithelia of vertebrates. *Micro. Res. Tech.* 23, 1-21.

- 30 Farbman, A.I. (1992). Cell biology of olfaction. In *Developmental and Cell Biology Series* (Cambridge University Press, Canada).

- 35 Garcia-Velasco, J. and Mondragon M. (1991). The incidence of the vomeronasal organ in 1000 human subjects and its possible clinical significance. *J. Steroid Biochem. Molec. Biol.* 39(4B), 561-564.

Henzel, W.J., Rodriguez, H., Singer, A.G., Stults, J.T.,
Macrides, F., Agosta, W.C., and Niall, H. (1988). The
primary structure of aphrodisin. J. Biol. Chem. 263(32),
16682-16687.

Jacobson, L. (1811). Description anatomique d'un organe
observe dans les Mammiferes. Ann. Mus. Hist Natn Paris
18, 412-424.

Jones, D.T., and Reed, R.R. (1989). Golf: An olfactory neuron specific G-protein involved in odorant signal transduction. *Science* 244, 790-795.

Kauer, J.S., Senseman, D.M., and Cohen, L.B. (1987).
Odor-elicited activity monitored simultaneously from 124
regions of the salamander olfactory bulb using a voltage-
sensitive dye. *Brain Res.* 418, 255-261.

- Keverne, E. B. (1983) Pheromonal influences on the endocrine regulation of reproduction. Trends Neurosc. 6 381-384.
- 5 Kevetter, G.A., and Winans, S.S. (1981). Connections of the corticomedial amygdala in the golden hamster. I. Efferents of the "Vomeronasal Amygdala." J. Comp. Neurol. 197, 81-98.
- 10 Krettek, J.E., and Price, J.L. (1977). Projections from the amygdaloid complex to the cerebral cortex and thalamus in the rat and cat. J. Comp. Neurol. 172, 687-722.
- 15 Krettek, J.E., and Price, J.L. (1978). Amygdaloid projections to subcortical structures within the basal forebrain and brainstem in the rat and cat. J. Comp. Neurol. 178, 225-254.
- 20 Lancet, D., Greer, C.A., Kauer, J.S., and Shepherd, G.M. (1982). Mapping of odor-related neuronal activity in the olfactory bulb by high resolution 2-deoxyglucose autoradiography. Proc. Natl. Acad. Sci. USA 79, 670-674.
- 25 Lomas, D.E., and Keverne, E.B. (1982). Role of the vomeronasal organ and prolactin in the acceleration of puberty in female mice. J. Reprod. Fert. 66, 101-107.
- 30 Ludwig, J., Marglit, T., Eismann, E., Lancet, D., and Kaupp, U.B. (1990). Primary structure of a cAMP-gated channel from bovine olfactory epithelium. FEBS Lett. 270, 24-29.
- 35 Macrides, F., Schoenfeld, T.A., Marchand, J.E., and Clancy, A. N. (1985). Evidence for morphologically,

neurochemically and functionally heterogeneous classes of mitral and tufted cells in the olfactory bulb. *Chemical Senses* 10(2), 175-202.

- 5 Melrose, D.R., Reed, H.C., and Patterson, R.L. (1971). Androgen steroids associated with boar odour as an aid to the detection of oestrus in pig artificial insemination. *Brit. Vet. J.* 127(10), 497-502.
- 10 Meredith, M. (1986). Vomeronasal organ removal before sexual experience impairs male hamster mating behavior. *Physiol. and Behavior* 36, 737-743.
- Meredith, M., and Fernandez-Fewell, G. (1994).
15 Vomeronasal system, LHRH, and sex behaviour. *Psychoneuroendocrinology*. 19 (5-7), 657-672.
- Michael, R.P., and Keverne, E.B. (1968). Pheromones in the communication of sexual status in primates. *Nature*
20 218(143), 746-749.
- Monti-Bloch L., Jennings-White, C., Dolberg, D. S., and Berliner, D. L. (1994). The human vomeronasal system. *Psychoneuroendocrinology*. 19 (5-7), 673-686.
25
- Moran, D.T, Jafek, B.W., and Rowley III, J. C. (1991). The vomeronasal (Jacobson's) organ in man: ultrastructure and frequency of occurrence. *J. Steroid Biochem. Molec. Biol.* 39(4B), 545-552.
30
- Mori, K., Mataga, N., and Imamura, K. (1992). Differential specificities of single mitral cells in rabbit olfactory bulb for a homologous series of fatty acid odor molecules. *J. Neurophysiol.* 67, 786-789.
35

Rogers, R.E., Dasgupta, P., Gubler, U., Grillo, M., Khew-Goodall, Y., and Margolis, F.L. (1987). Molecular cloning and sequencing of a cDNA for olfactory marker protein. Proc. Natl. Acad. Sci. USA 84, 1704-1708.

5

Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). Molecular Cloning: A Laboratory Manual, Second Edition (Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press).

10

Scalia, F., and Winans, S.S. (1975). The differential projections of the olfactory bulb and accessory olfactory bulb in mammals. J. Comp. Neur. 161, 31-56.

15

Schaeren-Wiemers, N., and Gerlin-Moser, A. (1993). A single protocol to detect transcripts of various types and expression levels in neural tissue and cultures cells: *in situ* hybridization using digoxigenin-labeled cRNA probes. Histochemistry 100, 431-440.

20

Singer, A.G. (1991). A chemistry of mammalian pheromones. J. Steroid Biochem. Molec. Biol. 39(4B), 627-632.

25

Sorensen, P.W., Hara, T.J., Stacey, N.E., and Goetz, F.W. (1988). F prostaglandins function as potent olfactory stimulants that comprise the postovulatory female sex pheromone in goldfish. Biol. of Reproduction 39, 1039-1050.

30

Stacey, N.E., and Sorensen, P.W. (1986). 17a, 20b-dihydroxy-4-pregnen-3-one: a steroidal primer pheromone increasing milt volume in the goldfish, *Carassius auratus*. Can. J. Zool. 64, 2412-2417.


35

- Stensaas, L.J., Lavker, R.M., Monti-Bloch, L., Grossier, B.I., and Berliner, D.L. (1991). Ultrastructure of the human vomeronasal organ. *J. Steroid Biochem. Molec. Biol.* 39(4B), 553-560.
- 5
- Stewart, W.B., Kauer, J.S., and Shepherd, G.M. (1979). Functional organization of rat olfactory bulb analyzed by the 2-deoxyglucose method. *J. Comp. Neurol.* 185, 715-734.
- 10
- Strader, C.D., Fong, T.M., Tota, M.R., and Underwood, D. (1994). Structure and function of G protein-coupled receptors. *Annu. Rev. Biochem.* 63, 101-132.
- 15
- Takami, S., and Graziadei, P.P. (1991). Light microscopic Golgi study of mitral/tufted cells in the accessory olfactory bulb of the adult rat. *J. Comp. Neurol.* 311, 65-83.
- 20
- Troemel, E., Chou, J., Dwyer, N.D., Colbert, H.A., and Bargmann, C.I. (1995). Divergent seven transmembrane receptors are candidate chemosensory receptors in *C. elegans*. *Cell*, in press.
- 25
- Vassar, R., Ngai, J., and Axel, R. (1993). Spatial segregation of odorant receptor expression in the mammalian olfactory epithelium. *Cell* 74, 309-318.
- 30
- Vassar, R., Chao, S.K., Sitcheran, R., Nunez, J.M., Vosshall, L.B., and Axel, R. (1994). Topographic organization of sensory projections to the olfactory bulb. *Cell* 79, 981-991.
- 35
- Winans, S.S., and Scalia, F. (1970). Amygdaloid nucleus: New afferent input from the vomeronasal organ.

5

10

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

Page 54 to 62 of specifications are References

☒ Scanned copy is best available. Drawings.